

Мартыненко А.С., Пикуз С.А., Рязанцев С.Н., Скобелев И.Ю., Баурд К.Д., Бузс Н., Вусли Н., Фенг Д., Доэль Л., Дюреи Ф., Кодама Р., Ксин Л., Ланкастер К., Мак Кена П., Мёрфи К.Д., Пикуз Т.А., Спиндо К., Фаенов А.Я., Фарли Д., Чен Л.М.

Определение параметров пикосекундной лазерной плазмы на начальных этапах её формирования методами высокоразрешающей рентгеновской спектроскопии

- [1] Kiriyama H. et al. High-contrast high-intensity repetitive petawatt laser // Optics Letters. — 2018. — jun. — Vol. 43, no. 11. — P. 2595. — Access mode: <https://www.osapublishing.org/abstract.cfm?URI=ol-43-11-2595>.
- [2] Фортов В. Е. Физика высоких плотностей энергии. — ФИЗМАТЛИТ, 2013. — ISBN: 978-5-9221-1468-4.
- [3] Mourou Gerard A., Tajima Toshiki, Bulanov Sergei V. Optics in the relativistic regime // Reviews of Modern Physics. — 2006. — apr. — Vol. 78, no. 2. — P. 309–371. — Access mode: <https://link.aps.org/doi/10.1103/RevModPhys.78.309>.
- [4] Danson C.N. et al. Vulcan petawatt - An ultra-high-intensity interaction facility // Nuclear Fusion. — 2004. — dec. — Vol. 44, no. 12. — P. S239–S246. — Access mode: <http://stacks.iop.org/0029-5515/44/i=12/a=S15?key=crossref.ad7333baa871b101225abbdc34d72c73>.
- [5] Faenov A Ya et al. High-performance x-ray spectroscopic devices for plasma microsources investigations // Physica Scripta. — 1994. — oct. — Vol. 50, no. 4. — P. 333–338. — Access mode: <http://stacks.iop.org/1402-4896/50/i=4/a=003?key=crossref.9f042b7c13029f622f645cd1b2fea7b2>.
- [6] Skobelev I Y et al. X-ray absorption spectroscopy study of energy transport in foil targets heated by petawatt laser pulses // Photonics Research. — 2018. — apr. — Vol. 6, no. 4. — P. 234. — Access mode: <https://www.osapublishing.org/abstract.cfm?URI=prj-6-4-234>.
- [7] MacFarlane J. J., Golovkin I. E., Woodruff P. R. HELIOS-CR - A 1-D radiation-magnetohydrodynamics code with inline atomic kinetics modeling // Journal of Quantitative Spectroscopy and Radiative Transfer. — 2006.
- [8] SPECT3D - A multi-dimensional collisional-radiative code for generating diagnostic signatures based on hydrodynamics and PIC simulation output / J. J. MacFarlane, I. E. Golovkin, P. Wang et al. // High Energy Density Physics. — 2007. — Vol. 3, no. 1-2. — P. 181.
- [9] Riley D. et al. K α yields from Ti foils irradiated with ultrashort laser pulses // Physical Review E. — 2005. — jan. — Vol. 71, no. 1. — P. 016406. — Access mode: <https://link.aps.org/doi/10.1103/PhysRevE.71.016406>.
- [10] Effects of nonionizing prepulses in high-intensity laser-solid interactions / K. B. Wharton, C. D. Boley, A. M. Komashko et al. // Physical Review E. — 2001. — jul. — Vol. 64, no. 2. — P. 025401. — Access mode: <https://link.aps.org/doi/10.1103/PhysRevE.64.025401>.
- [11] K-alpha conversion efficiency measurements for X-ray scattering in inertial confinement fusion plasmas / A.L. Kritcher, P. Neumayer, M.K. Urry et al. // High Energy Density Physics. — 2007. — may. — Vol. 3, no. 1-2. — P. 156–162. — Access mode: <https://linkinghub.elsevier.com/retrieve/pii/S1574181807000249>.
- [12] Chen S. N. et al. Creation of hot dense matter in short-pulse laser-plasma interaction with tamped titanium foils // Physics of Plasmas. — 2007. — oct. — Vol. 14, no. 10. — P. 102701. — Access mode: <http://aip.scitation.org/doi/10.1063/1.2777118>.
- [13] Alkhimova M. A. et al. High resolution X-ray spectra of stainless steel foils irradiated by femtosecond laser pulses with ultra-relativistic intensities // Optics Express. — 2017. — nov. — Vol. 25, no. 23. — P. 29501. — Access mode: <https://www.osapublishing.org/abstract.cfm?URI=oe-25-23-29501>.
- [14] Oks E. et al. In-depth study of intra-Stark spectroscopy in the x-ray range in relativistic laser-plasma interactions // Journal of Physics B: Atomic, Molecular and Optical Physics. — 2017. — Vol. 50, no. 24. — P. 245006.
- [15] Martynenko A. S. et al. Effect of plastic coating on the density of plasma formed in Si foil targets irradiated by ultra high-contrast relativistic laser pulses // Physical Review E. — 2019.
- [16] Martynenko A. S., Skobelev I. Yu., Pikuz S. A. Possibility of estimating high-intensity-laser plasma parameters by modelling spectral line profiles in spatially and time-integrated X-ray emission // Applied Physics B. — 2019. — feb. — Vol. 125, no. 2. — P. 31. — Access mode: <http://link.springer.com/10.1007/s00340-019-7149-4>.